

- Aggenbach, CJ, Kooijman, AM, Fuijta, Y, Van der Hagen, H, Van Til, M, Cooper, D. [&] Jones, L (2016). Does atmospheric nitrogen deposition lead to greater nitrogen and carbon accumulation in coastal sand dunes) *Biological Conservation*, <http://dx.doi.org/10.1016/j.biocon.2016.12.007>
- Benthien, O, Bober, J, Castens, J [&] Stolter, C (2016). Seed dispersal capacity of sheep and goats in a near-coastal dry grassland habitat. *Basic and Applied Ecology*. doi:10.1016/j.baae.2016.03.006
- Besley, A, Vijver, MG, Behrens, P [&] Bosker, T (2016). A standardized method for sampling and extraction methods for quantifying microplastics in beach sand. *Marine Pollution Bulletin*, Bird, E (2016). *Coastal Cliffs: Morphology and Management*. Springer.
- Costa, I., Reimann, T., Tsukamoto, S., Ludwig, J., Lindhorst, S., Frechen, M., Haase, H.C., Betzler, C. 2012. Comparison of OSL ages from young dune sediments with a high-resolution independent age model. *Quaternary Geochronology*, 10, 16-23.
- De Bryun, U. 2010. Flechten der Ostfriesischen Inseln. Schriftenreihe Nationalpark Niedersächsisches Wattenmeer Band 12.
- De Groot, AV, Oost, AP, Veeneklaas, RM, Lammerts, EJ, van Duin, WE [&] van Wesenbeeck, BK (2016). Tales of island tails: biogeomorphic development and management of barrier islands. *Journal of Coastal Conservation*. doi:10.1007/s11852-016-0446-8
- de Schipper, MA, de Vries, S, Ruessink, G, de Zeeuw, RC, Rutten, J., van Gelder-Maas, C., [&] Stive, M. J. (2016). Initial spreading of a mega feeder nourishment: Observations of the Sand Engine pilot project. *Coastal Engineering* 111, 23-38.
- Dekiff, J.H.; Remy, D.; Klasmeier, J.; Fries, E. 2014. Occurrence and spatial distribution of microplastics in sediments from Norderney. *Environmental Pollution* 186, 248-256
- Denga, J., Zhang, W., Harff, J., Schneider, R., Dudzinska-Nowak, J., Terefenko, P., Giza, A., Furmanczyk, K. 2014. A numerical approach for approximating the historical morphology of wave-dominated coasts. A case study of the Pomeranian Bight, southern Baltic Sea. *Geomorphology* 204, 425-443.
- Eigner, J. 2014. Der Meerfenchel (*Crithmum maritimum*) auf Pellworm. *Kieler Notizen zur Pflanzenkunde* 40, 53-55.
- Eschenbrenner, S, Gemeinholzer, B [&] Wissemann, V (2016). Morphologische, zytologische und populationsgenetische Untersuchungen zu Vorkommen und Status von *Drosera*-Arten in Schleswig-Holstein. *BfN-Skripten* 436, 175-183.
- Folmer, A, Revier, H. [&] Cupido, T (2016). Sustainable tourism development and the world heritage status of the Wadden Sea: The case of Terschelling. *Research in Hospitality Management* 6, 45-50.
- Garthe, S, Schwemmer, P, Paiva, VH, Corman, AM, Fock, HO, Voigt, CC [&] Adler, S (2016). Terrestrial and marine foraging strategies of an opportunistic seabird species breeding in the Wadden Sea. *Plos one* 11(8), e0159630.
- Grootjans, A, Shahrudin, R, Van de Craats, A, Kooijman, A, Ostermeijer, G, Petersen J, ... [&] Stuyfzand, P (2016): Window of *Liparis loeselii* populations during vegetation succession on the Wadden Sea islands. *Journal of Coastal Conservation*.
- Haacks, M, Janinhoff, N, Petersen, J, Stock, M, Dauck, H-P, Bertram, G (2016). Floristische Besonderheiten der Nordseeküstendünen Schleswig-Holsteins 2012-2014. *Kieler Notizen zur Pflanzenkunde* 41, 105–123.
- Heinecke, C. 2011. 2008-2010 auf den Ostfriesischen Inseln nachgewiesene Großschmetterlinge (*Lepidoptera*). *Drosera*, 2010, 117-132.
- Hellwig, U., Stock, M. (Eds.) 2014. *Dynamic Islands in the Wadden Sea*. Ecosystem No. 33: 1-134. Common Wadden Sea Secretariat, Wilhelmshaven, Germany. www.waddensea-secretariat.org, Wilhelmshaven Germany. http://www.waddensea-secretariat.org/news-and-service/news/dynamic_islands
- Herrling, G., C. Winter 2014. Morphological and sedimentological response of a mixed-energy barrier island tidal inlet to storm and fair-weather conditions. *Earth Surf. Dynam.* 2, 363-382.

- Hofstede, JL [&] Stock, M (2016). Climate change adaptation in the Schleswig-Holstein sector of the Wadden Sea: an integrated state governmental strategy. *Journal of Coastal Conservation*. doi:10.1007/s11852-016-0433-0
- Igel, J., Günther, T., Kuntzer, M. 2013. Ground-penetrating radar into coastal aquifer: the freshwater lens of Borkum Island. *Hydrology and Earth System Sciences*, 17, 519-531.
- Isermann, M., Rooney, P. 2014. Biological Flora of the British Isles: *Eryngium maritimum*. *Journal of Ecology* 102, 789-821.
- Jin, H.F., Liebezeit, G. 2013. Distribution of total mercury in coastal sediments from Jade Bay and its catchment, Lower Saxony, Germany. *Journal of Soils and Sediments*, 13, 441-449.
- Junghans, T. 2011. Auswirkungen verschiedener Straucharten auf die Artenvielfalt von Dünengebüsch auf Borkum (Ostfriesische Inseln, Niedersachsen). *Drosera*, 2010, 49-53.
- Kalisch, D. 2012. Relevance of crowding effects in a coastal National Park in Germany: results from a case study on Hamburger Hallig. *Journal of Coastal Conservation*, 16, 531-541.
- Koffijberg, K, Frikke, J, Hälterlein, B, Reichert, G [&] Andretzke, H. (2016). Breeding birds in trouble: A framework for an action plan in the Wadden Sea. 24 S.
- Kooijman, AM, Bruin, CJW., van de Craats, A, Grootjans, AP, Oostermeijer, JGB, Scholten, R [&] Sharudin, R (2016). Past and future of the EU-habitat directive species *Liparis loeselii* in relation to landscape and habitat dynamics in SW- Texel, the Netherlands. *Science of The Total Environment* 568: 107-117.
- Kooijman, AM, Van Til, M, Noordijk, E, Remke, E [&] Kalbitz, K (2016). Nitrogen deposition and grass encroachment in calcareous and acidic Grey dunes (H1230) in NW-Europe. *Biological Conservation*. <http://dx.doi.org/10.1016/j.biocon.2016.08.009>
- Koperski, M. 2011. Bryologische Beobachtungen auf den Ostfriesischen Inseln und Mellum. *Drosera*, 2010, 133-148.
- Kösters, F., Winter, C. 2014. Exploring German Bight coastal morphodynamics based on modelled bed shear stress. *Geo-Marine Letters* 34, 21-36.
- Kruckenbergh, H. [&] Schulze-Diekhoff, M (2016). Die Bedeutung von Stacheldraht als Gefahr für Vögel in Deutschland. *Corax* 23, 75-85.
- Lindhorst, S., Betzler, C. & Hass, H.C. 2013. Large scale architecture of a stacked Holocene spit - the stratigraphy of northern Sylt (southern North Sea). *Zeitschrift der Deutschen Gesellschaft für Geowissenschaften*, 164, 63-79.
- Mantilla-Contreras, J., Schirmel, J., Zerbe, S. 2012. Influence of soil and microclimate on species composition and grass encroachment in heath succession. *Journal of Plant Ecology*, 5, 249-259.
- Metzing, D., Kuhbier, H., Küver, B. 2011. *Crassula tillaea* (Crassulaceae) auf Baltrum - Erstnachweis für Niedersachsen. *Drosera*, 2010, 71-76.
- Petersen, J., Kers, B., Stock, M. 2014. TMAP Typology of Coastal Vegetation in the Wadden Sea Area. *Wadden Sea Ecosystem* 32, 1-86. <http://www.waddensea-secretariat.org/management/publications/tmap-typology-of-coastal-vegetation-in-the-wadden-sea-area>
- Reimann, T., Lindhorst, S., Thomsen, K.J., Murray, A.S., Frechen, M. 2012. OSL dating of mixed coastal sediment (Sylt, German Bight, North Sea). *Quaternary Geochronology*, 11, 52-62.
- Reise, K. 2014. A natural history of the Wadden Sea - Riddled by contingencies.
- Riksen, MJ, Goossens, D, Huiskes, HP, Krol, J [&] Slim, PA (2016). Constructing notches in foredunes: Effect on sediment dynamics in the dune hinterland. *Geomorphology* 253: 340-352.
- Ringel, H. 2013. Vegetationskartierung der Boddenufer am Kleinen Jasmunder Bodden (Insel Rügen). *Botanischer Rundbrief Mecklenburg-Vorpommern*, 50, 147-179.
- Roper, T., Greskowiak, J., Freund, H. & Massmann, G. 2013. Freshwater lens formation below juvenile dunes on a barrier island (Spiekeroog, Northwest Germany). *Estuarine Coastal and Shelf Science*, 121, 40-50.
- Röper, T., Kröger, K.F., Meyer, H., Sültenfuss, J., Greskowiak, J., Massmann, G. 2012. Groundwater ages, recharge conditions and hydrological evolution of a barrier island freshwater lens (Spiekeroog, Northern Germany). *Journal of Hydrology*, 454-455, 173-186.

- Schirmel, J. & Buchholz, S. 2013. Invasive moss alters patterns in life-history traits and functional diversity of spiders and carabids. *Biological Invasions*, 15, 1089-1100.
- Schirmel, J., Fartmann, T. 2014. Coastal heathland succession influences butterfly community composition and threatens endangered butterfly species. *Journal of Insect Conservation* 18, 111-120
- Schirmel, J., Mantilla-Contreras, J., Gauger, D., Blindow, I. 2014. Carabid beetles as indicators for shrub encroachment in dry grasslands. *Ecological Indicators* 49, 76-82.
- Schulz, K, Mikhailyuk, T, Dreßler, M, Leinweber, P [&] Karsten, U (2016). Biological soil crusts from coastal dunes at the Baltic sea: cyanobacterial and algal biodiversity and related soil properties. *Microbial Ecology* 71: 178-193.
- Schwemmer, P, Weiel, S [&] Garthe, S (2016). A fundamental study revisited: Quantitative evidence for territory quality in oystercatchers (*Haematopus ostralegus*) using GPS data loggers. *Ecology and Evolution*. DOI: 10.1002/ece3.2581
- Seer, FK, Irmeler, U, Schrautzer, J (2016). Beaches under pressure – effects of land use on vegetation at the Baltic Sea coast. *Journal of Applied Vegetation Science*. DOI: 10.1111/avsc.12221.
- Skowronek, S, Ewald, M, Isermann, M, Van de Kerchove, R, Lenoir, J, Aerts, R,... [&] Roccini, D (2016): Mapping an invasive bryophyte species using hyperspectral remote sensing data. *Biological Invasions*,
- Slob, AF, Geerdink, TRA, Röckmann, C, [&] Vöge, S 2016. Governance of the Wadden Sea. *Marine Policy* 71: 325-333.
- Sprichardt, J. 2011. Bienen und Wespen naturnaher Restheiden im Raum Cuxhaven. *Drosera*, 2010, 77-102.
- Stock, M. 2013 Der Schobüller Strand und seine Pflanzenwelt - früher und heute. *Kieler Notizen zur Pflanzenkunde* 39, 40-53.
- Stock, M., Baer, J. 2014. Trischen-wax and wane of a Wadden Sea island.
- Stuyfzand, PJ (2016). Observations and analytical modeling of freshwater and rainwater lenses in coastal dune systems. *Journal of Coastal Conservation*,
- Van der Zee, FF, de Knecht, B, Meeuwssen, HAM, Sanders, ME, Veraart, JA, Grashof-Bokdam, CJ, [&] Wegman, RMA (2016). Waterwinning en natuur (No. 2719). Alterra Wageningen UR.
- Walz, U., Stein, C. 2014. Indicators of hemeroby for the monitoring of landscapes in Germany. *Journal for Nature Conservation*. <http://dx.doi.org/10.1016/j.jnc.2014.01.007>
- Wasserstrom, H [&] Steinberger, Y (2016). Does distance from the sea affect a soil microarthropod community? *Acta Oecologica* 76: 39-46.
- Wegner, H. 2011. Die Kleinschmetterlings-Fauna ausgewählter Biotope auf der nordfriesischen Insel Sylt (Lepidoptera). *Drosera*, 2010, 1-44.
- Witt, R (2016). Vorkommen und Bestandssituation seltener Hummelarten (*Bombus*) in Niedersachsen und Bremen (Hymenoptera: Apidae). *Ampulex* 8.
- Wolff, W. J. 2013. Ecology of the Wadden Sea: Research in the past and challenges for the future. *Journal of Sea Research* 82, 3-9.
- Wünsch, Y., Schirmel, J., Fartmann, T. 2012. Conservation management of coastal dunes for Orthoptera has to consider oviposition and nymphal preferences. *Journal of Insect Conservation*, 16, 501-510.
- Zhang, W., Harff, J., Schneider, R., Meyer, M., Zorita, E., Hünicke, B. 2014. Holocene morphogenesis at the southern Baltic Sea: Simulation of multi-scale processes and their interactions for the Darss–Zingst peninsula. *Journal of Marine Systems* 129, 4-18